

Representation Formulas for Wave Equations with Profile Functions

Ben Schweizer

March 15, 2022 — PD22 virtual conference

- Part 1: Three wave equations, long-time-dispersion
 - Wave equation with constant coefficient
 - Wave equation in periodic media
 - Wave equation on a lattice
- Part 2: Formulas for profiles



Wave equation with constant coefficient

Wave equation

$$\partial_t^2 u(x, t) = c^2 \Delta u(x, t)$$

Coefficient $a = c^2$ independent of x



Planar-wave ansatz: $u(x, t) = e^{ik \cdot x} e^{-i\omega t}$

u is a solution if and only if $\omega^2 = c^2 |k|^2$

Result: The dispersion relation is $\omega(k) = \pm c|k|$

A solution:

$$u(x, t) = \int_{\mathbb{R}^d} u_0(k) e^{ik \cdot x} e^{-ic|k|t} dk$$


Satisfy initial conditions by using $+$ and $-$

One dimensional equation: An exact solution is given by shifts,

$$u(x, t) = \frac{1}{2} u_0(x - ct) + \frac{1}{2} u_0(x + ct)$$

Compare Fourier representation: Solutions depend on $x + ct$ and $x - ct$

Wave equation in periodic media



Let $a : \mathbb{R}^d \rightarrow (\delta, \infty)$ be 1-periodic
Set $a_\varepsilon(x) := a(x/\varepsilon)$

Homogenization problem


$$\partial_t^2 u^\varepsilon = \nabla \cdot (a_\varepsilon(x) \nabla u^\varepsilon(x))$$

Classical homogenization: $u^\varepsilon \approx u$, where u solves the


Homogenized equation

$$\partial_t^2 u(x, t) = \nabla \cdot a_* \nabla u(x, t)$$


Observation in dimension 1 with periodicity $\varepsilon = 1/6$, the solution of homogenized equation is a shift, $u(x, t) = u_0(x - t)$



Time $t = 10$



Time $t = 50$



Time $t = 120$

→ We observe dispersion for large times!

Result on weak dispersion

Theorem (T.Dohnal, A.Lamacz, B.S., 2014 and 2015)


There exist $A, E \in \mathbb{R}^{d \times d}$ and $F \in \mathbb{R}^{d \times d \times d \times d}$ such that:

Solutions of the well posed **weakly dispersive** effective equation


$$\partial_t^2 w^\varepsilon = AD^2 w^\varepsilon + \varepsilon^2 ED^2 \partial_t^2 w^\varepsilon - \varepsilon^2 FD^4 w^\varepsilon$$

satisfy the approximation estimate

$$\sup_{t \in [0, T\varepsilon^{-2}]} \|u^\varepsilon(\cdot, t) - w^\varepsilon(\cdot, t)\|_{L^2(\mathbb{R}^d) + L^\infty(\mathbb{R}^d)} \leq C_0 \varepsilon$$



u^ε : solution for coefficient a_ε



w^ε : solution of weakly dispersive equation



The proof is based on an explicit formula for solutions:

Proposition (Bloch-wave approximation of u^ε)

$$v^\varepsilon(x, t) := \frac{1}{2\sqrt{2\pi}^n} \sum_{\pm} \int_K F_0(k) e^{ik \cdot x} \exp\left(\pm it \sqrt{\sum A_{lm} k_l k_m}\right) \\ \times \exp\left(\pm \frac{i\varepsilon^2}{2} t \frac{\sum C_{lmnq} k_l k_m k_n k_q}{\sqrt{\sum A_{lm} k_l k_m}}\right) dk$$
$$\sup_{t \in [0, T\varepsilon^{-2}]} \|u^\varepsilon(\cdot, t) - v^\varepsilon(\cdot, t)\|_{(L^2 + L^\infty)(\mathbb{R}^n)} \leq C_0 \varepsilon$$

- Taylor expansion of the smallest Bloch eigenvalue $\mu_0(k)$
- Replacement trick to re-write the equation
- Estimate for $u^\varepsilon - v^\varepsilon$ from approximations in the explicit formula
- Estimate for $v^\varepsilon - w^\varepsilon$ by energy methods

Waves in a lattice



Periodicity of the lattice: $\varepsilon > 0$
Dimension $d \geq 1$, lattice points
 $\gamma \in \varepsilon\mathbb{Z}^d$
Displacement at time $t \in [0, \infty)$
is $u^\varepsilon(\gamma, t)$.

Evolution of displacements

Wave equation in lattice:

$$\partial_t^2 u^\varepsilon(\gamma, t) = \frac{1}{\varepsilon^2} \sum_{j \in \mathbb{Z}^d} a_j u^\varepsilon(\gamma + \varepsilon j, t)$$

Initial conditions: $u^\varepsilon(\gamma, 0) = u_0(\gamma)$ and $\partial_t u^\varepsilon(\gamma, 0) = u_1(\gamma)$.

Example: Dimension $d = 1$, $a_1 = a_{-1} = 1$, $a_0 = -2$, and $a_j = 0$ else.

Explicit solutions for lattice waves in Fourier space

The equation for the **Fourier transform**

$$\hat{u}^\varepsilon(k, t) = \varepsilon^d \sum_{\gamma \in (\varepsilon\mathbb{Z})^d} e^{-ik \cdot \gamma} u^\varepsilon(\gamma, t)$$

is easily calculated: For every $k \in \mathbb{R}^d$ holds

$$\varepsilon^2 \partial_t^2 \hat{u}^\varepsilon(k, t) = -\omega(\varepsilon k)^2 \hat{u}^\varepsilon(k, t) \quad \text{with} \quad \omega(k)^2 := - \sum_{j \in \mathbb{Z}^d} a_j e^{ik \cdot j}$$

The **dispersion relation** in the example is $\frac{\omega(\varepsilon k)}{\varepsilon} = \sqrt{k^2 - \frac{1}{12}\varepsilon^2 k^4} \pm \dots$

Explicit solution in Fourier space

$$\hat{u}^\varepsilon(k, t) \sim e^{\pm i[\omega(\varepsilon k)/\varepsilon]t}$$

Also for lattices there is a weakly dispersive limit equation for w^ε

Theorem (B.S. & F.Theil, 2018)

$$\sup_{t \leq T/\varepsilon^2} \|\partial_t \hat{w}^\varepsilon(\cdot, t) - \partial_t \hat{u}^\varepsilon(\cdot, t)\|_{L^2(\mathbb{R}^d)} \leq C\varepsilon$$

Part 2: Profile equations

Wave in heterogeneous media

$$\partial_t^2 u^\varepsilon = \nabla \cdot (a_\varepsilon(x) \nabla u^\varepsilon(x))$$

Weakly dispersive wave equation

$$\partial_t^2 w^\varepsilon = AD^2 w^\varepsilon + \varepsilon^2 ED^2 \partial_t^2 w^\varepsilon$$

Fourier representation

$$\hat{u}^\varepsilon(k, t) \sim e^{\pm i[\omega(\varepsilon k)/\varepsilon]t}$$

Profile equation

$$\partial_\tau V(z, \tau; q) = b(q) \partial_z^3 V$$




Profile in 1D: Linearized KdV equation

Task: Study the long time behavior of

$$\partial_t^2 u^\varepsilon = c^2 \partial_x^2 u^\varepsilon + \varepsilon^2 B \partial_x^4 u^\varepsilon$$

We want an equation for *this* \rightarrow



Solution at time $t = 120$

Ansatz $u^\varepsilon(x, t) = V(x - ct, \varepsilon^2 t)$. Inserting yields

$$c^2 \partial_z^2 V - 2c \varepsilon^2 \partial_z \partial_\tau V + O(\varepsilon^4) = c^2 \partial_z^2 V + \varepsilon^2 B \partial_z \partial_z^3 V$$

Profile evolution equation: Linearized KdV equation; direction $q = \pm 1$:

$$\partial_\tau V^\varepsilon(z, \tau; q) = b(q) \partial_z^3 V^\varepsilon(z, \tau; q)$$

Initial condition:

$$\hat{V}_0^\varepsilon(\xi; \pm 1) := \begin{cases} \hat{u}_0^\varepsilon(\pm \xi) & \text{for } \xi > 0 \\ 0 & \text{else} \end{cases}$$



Reconstruction:

$$v^\varepsilon(x, t) := \begin{cases} V^\varepsilon(|x| - ct, \varepsilon^2 t; +1) & \text{for } x > 0 \\ V^\varepsilon(|x| - ct, \varepsilon^2 t; -1) & \text{for } x < 0 \end{cases}$$

2-dimensional solution

Theorem (Lattice solution and profile solution)

$$\|\hat{u}^\varepsilon(\cdot, \tau/\varepsilon^2) - \hat{v}^\varepsilon(\cdot, \tau/\varepsilon^2)\|_{L^2(\mathbb{R})} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$



The profile is given by a linearized KdV equation

$$\partial_\tau V^\varepsilon(z, \tau; q) = b(q) \partial_z^3 V^\varepsilon(z, \tau; q)$$

Initial condition

$$\hat{V}_0^\varepsilon(\xi; q) = \begin{cases} \sqrt{\frac{\xi}{2\pi i}} \hat{u}_0^\varepsilon(\xi q) & \text{for } \xi > 0 \\ 0 & \text{else} \end{cases}$$

Reconstruction

$$v^\varepsilon(x, t) = \frac{1}{|x|^{1/2}} V^\varepsilon \left(|x| - ct, \varepsilon^2 t; \frac{x}{|x|} \right)$$

General functions in Fourier space

- Wave equation

$$\hat{u}^\varepsilon(k, t) = \hat{u}_0(k) e^{-ic|k|t}$$

- Lattice

$$\hat{u}^\varepsilon(k, t) = \hat{u}_0(k) e^{-ic|k|t} e^{i\frac{b}{2}|k|^3\varepsilon^2 t}$$

- Periodic media

$$\hat{u}^\varepsilon(k, t) = \hat{u}_0(k) e^{-i|k|A t} e^{ib(k)\varepsilon^2 t}$$

Main result:

multiplication $\hat{u}_0(k) \mapsto e^{ic|k|\tau/\varepsilon^2} e^{-ib(|k|)\tau} \hat{u}_0(k)$

is like application of $\hat{Q}_b^\rho = \mathcal{F}_d \circ \mathcal{S} \circ \mathcal{F}_1^{-1} \circ J_b \circ \mathcal{R}_\rho$

Theorem (A.Lamacz & B.S., 2020)

Let u_0 be initial data, $d \leq 3$, $b : \mathbb{R} \rightarrow \mathbb{R}$ a dispersion function (appropriate). For $\tau > 0$ and $k \in \mathbb{R}^d$ with $|k| > \rho$, as $\varepsilon \rightarrow 0$:

$$(\hat{Q}_b^\rho \hat{u}_0)(k, \tau/\varepsilon^2) - e^{ic|k|\tau/\varepsilon^2} e^{-ib(|k|)\tau} \hat{u}_0(k) \rightarrow 0$$

Abstract reconstruction operators

Dimension $d \in \{1, 2, 3\}$, direction $q \in S^{d-1} \subset \mathbb{R}^d$

Restriction operator \mathcal{R} : For $u_0 = u_0(x)$

$$\hat{V}_0(\xi; q) := (\mathcal{R}\hat{u}_0)(\xi; q) := \left(\frac{|\xi|}{2\pi i}\right)^{(d-1)/2} \mathbf{1}_{\{\xi>0\}} \hat{u}_0(\xi q)$$

Profile variables: ξ dual to z

Multiplication operator J_b : For $b : \mathbb{R} \rightarrow \mathbb{R}$ set


$$\hat{V}(\xi, \tau; q) := (J_b \hat{V}_0)(\xi, \tau; q) := e^{-ib(\xi)\tau} \hat{V}_0(\xi; q)$$

Shell operator \mathcal{S} : For a profile $V = V(z, \tau; q)$


$$(SV)(x, t) := \frac{1}{(ct)^{(d-1)/2}} \mathbf{1}_{\{|x|<2ct\}} V\left(|x| - ct, \varepsilon^2 t; \frac{x}{|x|}\right)$$

The reconstruction operator \mathcal{Q}_b :


$$\mathcal{Q}_b = \mathcal{S} \circ \mathcal{F}_1^{-1} \circ J_b \circ \mathcal{R}$$



Cut of u_0 and of \hat{u}_0



$V_0(\cdot; q)$ and $V(\cdot, \tau; q)$




Stationary phase

Key in the proof: Write the reconstruction

$$v(x, t) := \frac{1}{(ct)^{(d-1)/2}} \mathbf{1}_{\{|x| < 2ct\}} V \left(|x| - ct, \varepsilon^2 t; \frac{x}{|x|} \right)$$

in Fourier space



Lemma (Stationary phase)

$d \in \{1, 2, 3\}$ and $\kappa \in S^{d-1}$ a point. Test-function $\phi \in C^1(S^{d-1}; \mathbb{R})$ supported in half sphere $\{q \in S^{d-1} \mid q \cdot \kappa \geq 0\}$. Then, as $N \rightarrow \infty$:

$$(2\pi i)^{-(d-1)/2} \int_{S^{d-1}} N^{(d-1)/2} e^{i(1-q \cdot \kappa)N} \phi(q) dS(q) \longrightarrow \phi(\kappa)$$

- **Dispersion:** Can appear in ε -heterogeneous wave equations at time scale $t \sim \varepsilon^{-2}$
A one-dimensional limit equation can be $\partial_t^2 w^\varepsilon = \partial_x^2 w^\varepsilon + \varepsilon^2 \partial_x^2 \partial_t^2 w^\varepsilon$
- **Profile equation:** Linearized KdV equation $\partial_\tau V = b \partial_z^3 V$
- **Abstract reconstruction:** Quite general evolutions in Fourier space can be approximated with profiles

Thank you!

- [1] A.Lamacz: Dispersive effective models for waves in heterogeneous media. *Math. Models Methods Appl. Sci.* 21(9), 2011
- [2] T.Dohnal, A.Lamacz, B.Schweizer: Bloch-wave homogenization on large time scales and dispersive effective wave equations. *Multiscale Model. Simul.* 12(2), 2014
- [3] T.Dohnal, A.Lamacz, B.Schweizer: Dispersive homogenized models and coefficient formulas for waves in general periodic media. *Asymptot. Anal.* 93(1-2), 2015
- [4] B.Schweizer, F.Theil: Lattice dynamics on large time scales and dispersive effective equations, *SIAM J. Appl. Math.* 78(6), 2018
- [5] A.Lamacz, B.Schweizer: Representation of solutions to wave equations with profile functions, *Analysis and Applications*, 2020